

DIRAC OSCILLATOR VIA R-DEFORMED HEISENBERG ALGEBRA

R. de Lima Rodrigues¹

Centro Brasileiro de Pesquisas Físicas (CBPF)

Rua Dr. Xavier Sigaud, 150, CEP 22290-180, Rio de Janeiro-RJ, Brazil

A. N. Vaidya

Instituto de Física - Universidade Federal do Rio de Janeiro

Caixa Postal 68528 - CEP 21945-970, Rio de Janeiro, Brazil

Abstract

The complete energy spectrum for the Dirac oscillator via R-deformed Heisenberg algebra is investigated.

PACS numbers: 11.30.Pb, 03.65.Fd, 11.10.Ef.

To appear in the proceedings at the XXIII Brazilian National Meeting on Particles and Fields, October 15 to 19, 2002, Águas de Lindóia-SP, Brazil, site www.sbf1.if.usp.br/eventos/enfpc/xxiii.

1 Introduction

The relativistic Dirac oscillator proposed by Moshinsky-Szczepaniak [1] is a spin $\frac{1}{2}$ object with the Hamiltonian which in the non-relativistic limit leads to that of a 3-dimensional isotropic oscillator shifted by a constant term plus a $\vec{L} \cdot \vec{S}$ coupling term for both signs of energy. There they construct a Dirac Hamiltonian, linear in the momentum $\vec{p}$ and position $\vec{r}$, whose square leads to the ordinary harmonic oscillator in the non-relativistic limit. The Dirac oscillator have been investigated in several context [2].

The R-deformed Heisenberg algebra or Wigner-Heisenberg algebraic technique [3] was recently super-realized for the SUSY isotonic oscillator [4, 5]. The R-Heisenberg algebra has also been investigated for the three-dimensional non-canonical oscillator to generate a representation of the orthosymplectic Lie superalgebra $osp(3/2)$ [6].

¹Permanent address: Departamento de Ciências Exatas e da Natureza, Universidade Federal de Campina Grande, Cajazeiras - PB, 58.900-000-Brazil. E-mail to RLR is rafaelr@cbpf.br or rafael@fisica.ufpb.br, the e-mail to ANV is vaidya@if.ufrj.br.

The R-Heisenberg algebra has been found relevant in the context of integrable models [7], and the Calogero interaction [8, 9]. Recently it has been employed for bosonization of supersymmetry in quantum mechanics [10], and the discrete space structure for the 3D Wigner quantum oscillator has been investigated [12]. In this work, we obtain the complete energy spectrum for the Dirac oscillator via R-deformed Heisenberg (RDH) algebra.

2 3D Wigner Oscillator

In this Section, we provide a three dimension presentation of the Wigner system with its bosonic sector to be the 3D isotropic oscillator (assumed to be of spin- $\frac{1}{2}$, to aid factorization).

The R-deformed Heisenberg (or Wigner-Heisenberg) algebra is given by following (anti-)commutation relations ($[A, B]_+ \equiv AB + BA$ and $[A, B]_- \equiv AB - BA$) :

$$H = \frac{1}{2}[a^-, a^+]_+, \quad [H, a^\pm]_- = \pm a^\pm, \quad [a^-, a^+]_- = 1 + cR, \quad [R, a^\pm]_+ = 0, \quad R^2 = 1, \quad (1)$$

where c is a real constant associated to the Wigner parameter [4]. Note that when $c = 0$ we have the standard Heisenberg algebra.

It is straightforward, following the analogy with the Ref. [4], to define the super-realizations for the ladder operators $a^\mp(\vec{\sigma} \cdot \vec{L} + \mathbf{1})$ for $H_W \equiv H(\vec{\sigma} \cdot \vec{L} + \mathbf{1})$ taking the explicitly forms

$$a^\mp = a^\mp(\vec{\sigma} \cdot \vec{L} + \mathbf{1}) = \frac{1}{\sqrt{2}} \left\{ \mp \Sigma_1 \left(\frac{\partial}{\partial r} + \frac{1}{r} \right) \pm \frac{1}{r} (\vec{\sigma} \cdot \vec{L} + \mathbf{1}) \Sigma_1 \Sigma_3 - \Sigma_1 r \right\} \quad (2)$$

which satisfy together with $H_W \equiv H(\vec{\sigma} \cdot \vec{L} + \mathbf{1})$ all the algebraic relations of the RDH algebra with the constant $\frac{c}{2}$ replaced by $(\vec{\sigma} \cdot \vec{L} + \mathbf{1})$ and $R = \Sigma_3$. Note that $(\vec{\sigma} \cdot \vec{L} + \mathbf{1})$ commutes with all the basic elements ($a^\mp$ and H_W) of the RDH algebra.

It may be observed that the RDH algebra that gets defined here is in fact three dimensional (one dimension for r and two for $(\vec{\sigma} \cdot \vec{L} + \mathbf{1})$) and is identically satisfied on any arbitrary three dimensional wave function.

On the eigenspaces of the operator $(\vec{\sigma} \cdot \vec{L} + \mathbf{1})$, the 3D Wigner algebra gets reduced to a 1D from with $(\vec{\sigma} \cdot \vec{L} + \mathbf{1})$ replaced by its eigenvalue $\mp(\ell + 1)$, $\ell = 0, 1, 2, \dots$, where ℓ is the orbital angular momentum quantum number. The eigenfuncitons of $(\vec{\sigma} \cdot \vec{L} + \mathbf{1})$ for the eigenvalues $(\ell + 1)$ and $-(\ell + 1)$ are respectivaly given by the well known spin-spherical harmonic $y_\mp$.

Now, considering simultaneous eigenfuncitons of the mutually commuting H_W and $(\vec{\sigma} \cdot \vec{L} + \mathbf{1})$ by

$$\psi_{W,+} = \begin{pmatrix} \tilde{R}_{1,+}(r) \\ \tilde{R}_{2,+}(r) \end{pmatrix} y_+, \quad (\vec{\sigma} \cdot \vec{L} + \mathbf{1})\psi_{W,+} = (\ell + 1)\psi_{W,+}, \quad (3)$$

$$\psi_{W,-} = \begin{pmatrix} \tilde{R}_{1,-}(r) \\ \tilde{R}_{2,-}(r) \end{pmatrix} y_-, \quad (\vec{\sigma} \cdot \vec{L} + \mathbf{1})\psi_{W,-} = -(\ell + 1)\psi_{W,-}, \quad (4)$$

(where the use of the subscript $+$ ($-$) indicates association with $[y_+(y_-)]$, we observe that the positive semi-definite form of H_W the ladder relations and the form of H_W dictat that the ground state energy $E_w^{(0)}(\vec{\sigma} \cdot \vec{L} + \mathbf{1}) \geq 0$, where $E_W(\vec{\sigma} \cdot \vec{L} + \mathbf{1})$ indicates a function of $\vec{\sigma} \cdot \vec{L} + \mathbf{1}$, is determined by the annihilation condition which reads as two cases.

3 The Dirac Oscillator Model

Adding an "anomalous momentum" in the form of a (nonlocal) linear and hermitian interaction, $\vec{\alpha} \cdot \vec{\pi} \equiv -iM\omega\beta\vec{\alpha} \cdot \vec{r} = (\vec{\alpha} \cdot \vec{\pi})^\dagger$, in the (noncovariant) Dirac free particle equation with mass M and spin- $\frac{1}{2}$, in the natural sistem of units,

$$i\frac{\partial\psi}{\partial t} = (\vec{\alpha} \cdot \vec{p} + M\beta)\psi, \quad (5)$$

one obtains the equation for the Dirac oscillator [1]:

$$i\frac{\partial\psi}{\partial t} = \{\vec{\alpha} \cdot (\vec{p} + \vec{\pi}) + M\beta\}\psi, \quad (6)$$

where M and ω are, respectively, the mass of the particle and the frequency of the oscillator, and the matrices $(\vec{\alpha}, \beta)$ satisfy the following properties:

$$[\alpha_i, \beta]_+ = 0, \quad [\alpha_i, \alpha_j]_+ = 2\delta_{ij}\mathbf{1}, \quad \beta^2 = \mathbf{1} = \alpha_i^2, \quad (i, j = 1, 2, 3). \quad (7)$$

Writing the Dirac spinor in terms of the upper and lower components, respectively, ψ_1 and ψ_2 , $\Psi(\vec{r}, t) = \exp(-iEt) \begin{bmatrix} \psi_1(\vec{r}) \\ \psi_2(\vec{r}) \end{bmatrix}$ the standard representation of the matrices $\vec{\alpha}$ and β .

4 The Dirac oscillator via RDH algebra

In this section, we implement a new realization of the Dirac oscillator in terms of elements of the R-deformed Heisenberg algebra. To solve the equation Dirac, following the usual procedure, we consider the second order differential equation,

$$\tilde{H}_D\psi(\vec{r}) = E\psi(\vec{r}), \quad (8)$$

where $\tilde{H}_D$ is a second order Hamiltonian, $\tilde{H}_D = H_D^2 + M^2 \mathbf{1}$, $\tilde{E} = \frac{E^2 - M^2}{2M}$. In the spherical polar coordinate system, we obtain the non-relativistic form of the Hamiltonian U [11], for an isotropic 3D SUSY harmonic oscillator with spin- $\frac{1}{2}$.

We consider a unitary operator in terms of the radial projection of the spin,

$$U = \begin{bmatrix} 1 & 0 \\ 0 & \sigma_r \end{bmatrix} = U^{-1} = U^\dagger, \quad (9)$$

to obtain the following relation between the transformed Dirac Hamiltonian, $\tilde{H}_D$, the 3D Wigner Hamiltonian, H_W , and the SUSY Hamiltonian, H_{SUSY} [11]:

$$H_{\text{SUSY}} = U \tilde{H}_D U^\dagger = H_W - \frac{1}{2} \{1 + 2(\boldsymbol{\sigma} \cdot \vec{L} + \mathbf{1}) \Sigma_3\} \omega \Sigma_3. \quad (10)$$

4.1 The energy spectrum of the Dirac oscillator

The energy spectra of the operators $\tilde{H}_D$ and H_{SUSY} are identical, since these operators are related by a unitary transformation. However, the relation between the principal quantum number N and the angular momentum (ℓ) is different, in each case. Obviously, the energy spectrum associated with the two types of eigenspaces belonging to the eigenvalues $\pm(\ell + 1)$:

$$\text{Case(i)} \rightarrow \vec{\sigma} \cdot \vec{L} + \mathbf{1} \rightarrow \ell + 1 = j + \frac{1}{2}, \quad j = \ell + \frac{1}{2}$$

$$\tilde{E}_{N\ell} = \frac{E^2 - M^2}{2M} = \begin{cases} 2m\omega = \tilde{E}_{N(\ell+1)}^+, \\ 2(m+1)\omega = \tilde{E}_{N\ell}^-, \end{cases} \quad (11)$$

where $m = 0, 1, 2, \dots$.

$$\text{Case(ii)} \rightarrow \vec{\sigma} \cdot \vec{L} + \mathbf{1} \rightarrow -(\ell + 1) = -(j + \frac{1}{2}), \quad j = (\ell + 1) - \frac{1}{2} :$$

$$\tilde{E}_{N\ell} = \frac{E^2 - M^2}{2M} = \begin{cases} (N + j + 3/2)\omega = \tilde{E}_{N\ell}^+, & N = j - \frac{1}{2}, j + 3/2, j + 7/2, \dots, \\ (N + j + 5/2)\omega = \tilde{E}_{N(\ell+1)}^-, & N = j + \frac{1}{2}, j + 5/2, \dots. \end{cases}$$

5 Conclusion

In this work we investigate the Dirac oscillator with the help of techniques of super-realization of the R-deformed Heisenberg algebra.

The Dirac oscillator with different interactions has been treated by Castaños *et al.* and by Dixit *et al.* [2]. These works motivate the construction of a new linear Hamiltonian in terms of the momentum, position and mass coordinates, through a set of seven mutually anticommuting 8x8-matrices yielding a representation of the Clifford algebra $\mathcal{Cl}_7$. The seven elements of the Clifford algebra $\mathcal{Cl}_7$ generate the three linear momentum components, the three position coordinates components and the mass, and their squares are the 8x8-identity matrix $\mathbf{I}_{8 \times 8}$. Results of our analysis on Dirac oscillator via the Clifford algebra $\mathcal{Cl}_7$ are in preparation.

In a forthcoming paper we show that the Dirac oscillator equation can be resolved algebraically without having to transform it into a second order differential equation. Therefore, the important connection for the Dirac 3D-isotropic oscillator with the linear ladder operators of the R-deformed Heisenberg algebra, satisfying the concomitant general oscillator quantum rule of Wigner, have explicit in this work.

ACKNOWLEDGMENTS

The authors are grateful to J. Jayaraman, whose advises and encouragement were precious. RLR was supported in part by CNPq (Brazilian Research Agency). He wishes to thank J. A. Helayel Neto for the kind of hospitality at CBPF-MCT. The authors wish also thank the staff of the CBPF and DCEN-CFP-UFCG. The authors are also grateful to the organizing committee of the of the XXIII Brazilian National Meeting on Particles and Fields (ENFPC), October 15 to 19, 2002, Águas de Lindóia-SP.

References

- [1] M. Moshinsky and A. Szczepaniak, *J. Phys. A: Math. Gen.* **22**, L817 (1989).
- [2] M. Moreno and A. Zentella, *J. Phys. A: Math. Gen.* **22**, L821 (1989); J. Beckers and N. Debergh, *Phys. Rev.* **D42**, 1255 (1990); C. Quesne and M. Moshinsky, *J. Phys. A: Math. Gen.* **23**, 2263 (1990); J. Benítez, R. P. Martínez y Romero, H. N. Núñez-Yépez and A. L. Salas-Brito, *Phys. Rev. Lett.* **64**, 1643 (1990); R. P. Martínez y Romero, Matías Moreno and A. Zentella, *Phys. Rev.* **D43**, 2036 (1991); O. L. de Lange, *J. Phys. A* **24**, 667 (1991); O. L. de Lange and R. E. Raab, *J. Math. Phys.* **32**, 1296 (1991); O. Castaños, A. Frank, R. López and L. F. Urrutia, *Phys. Rev.* **D43**, 544 (1991); V. V. Dixit, T. S. Santhanam, and W. D. Thacker, *J. Math. Phys.* **33**, 1114 (1992).
- [3] E. P. Wigner, *Phys. Rev.* **77**, 711 (1950); L. M. Yang, *Phys. Rev.* **84**, 788 (1951); L. O’Raifeartaigh and C. Ryan, *Proc. R. Irish Acad.* **A62**, 93 (1963); Y. Ohnuki and S. Kamefuchi, *J. Math. Phys.* **19**, 67 (1978); Y. Ohnuki and S. Watanabe, *J. Math. Phys.* **33**, 3653 (1992).
- [4] J. Jayaraman and R. de Lima Rodrigues, *J. Phys. A: Math. Gen.* **23**, 3123 (1990).
- [5] S. M. Plyushchay, *Int. J. Mod. Phys.* **A15**, 3679 (2000) and references therein.
- [6] T. D. Palev and N. I. Stoilova *J. Phys. A: Math. Gen.* **27**, 7387 (1994).
- [7] M. A. Vasiliev, *Int. J. Mod. Phys.* **A6**, 1115, (1991).

- [8] A. P. Polychronakos, *Phys. Rev. Lett.* **69**, 703 (1992).
- [9] T. Brzezinski, I. L. Egusquiza and A. J. Macfarlane, *Phys. Lett.* **B311**, 202 (1993); L. Brink, T. H. Hansson and M. A. Vasiliev, *Phys. Lett.* **B286**, 109 (1992); L. Brink, T. H. Hansson and S. Konstein and M. A. Vasiliev, *Nucl. Phys.* **B401**, 591 (1993).
- [10] S. M. Plyushchay, *Nucl. Phys.* **B491**, 619 (1997).
- [11] H. Ui, *Prog. Theor. Phys.* **72**, 813 (1984); H. Ui and G. Takeda, *Prog. Theor. Phys.* **72**, 266 (1984); A. B. Balantekin, *Ann. of Phys.* **164**, 277 (1985).
- [12] R. C. King, T. D. Palev, N. I. Stoloiva and J. Van der Jeugt, *Discrete space structure for the 3D Wigner quantum oscillator*, hep-th/0210164.